


Anti-mTOR antibody



Description	Rabbit polyclonal to mTOR.
Model	STJ94280
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IF, IHC-p, WB
Immunogen	Synthesized peptide derived from human mTOR around the non-phosphorylation site of S2448.
Immunogen Region	2390-2470 aa
Gene ID	2475
Gene Symbol	MTOR
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:40000
Specificity	mTOR polyclonal antibody detects endogenous levels of mTOR protein.
Tissue Specificity	Expressed in numerous tissues, with highest levels in testis.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	FOR RESEARCH USE ONLY (RUO).
Protein Name	Serine/Threonine-Protein Kinase MtorFk506-Binding Protein 12-Rapamycin Complex-Associated Protein 1Fkbp12-Rapamycin Complex-Associated ProteinMammalian Target Of RapamycinMtorMechanistic Target Of RapamycinRapamycin And Fkbp12 Target 1Rapamycin Target Protein 1

Molecular Weight	288 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:3942OMIM:601231
Alternative Names	Anti-Serine/Threonine-Protein Kinase Mtor antibodyAnti-Fk506-Binding Protein 12-Rapamycin Complex-Associated Protein 1 antibodyAnti-Fkbp12-Rapamycin Complex-Associated Protein antibodyAnti-Mammalian Target Of Rapamycin antibodyAnti-Mtor antibodyAnti-Mechanistic Target Of Rapamycin antibodyAnti-Rapamycin And Fkbp12 Target 1 antibodyAnti-Rapamycin Target Protein 1 antibodyAnti-MTOR FRAP FRAP1 FRAP2 RAFT1 RAP1 antibody
Function	<p>Serine/threonine protein kinase which is a central regulator of cellular metabolism, growth and survival in response to hormones, growth factors, nutrients, energy and stress signals. MTOR directly or indirectly regulates the phosphorylation of at least 800 proteins. Functions as part of 2 structurally and functionally distinct signaling complexes mTORC1 and mTORC2 (mTOR complex 1 and 2). Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. This includes phosphorylation of EIF4EBP1 and release of its inhibition toward the elongation initiation factor 4E (eIF4E). Moreover, phosphorylates and activates RPS6KB1 and RPS6KB2 that promote protein synthesis by modulating the activity of their downstream targets including ribosomal protein S6, eukaryotic translation initiation factor EIF4B, and the inhibitor of translation initiation PDCD4. Stimulates the pyrimidine biosynthesis pathway, both by acute regulation through RPS6KB1-mediated phosphorylation of the biosynthetic enzyme CAD, and delayed regulation, through transcriptional enhancement of the pentose phosphate pathway which produces 5-phosphoribosyl-1-pyrophosphate (PRPP), an allosteric activator of CAD at a later step in synthesis, this function is dependent on the mTORC1 complex. Regulates ribosome synthesis by activating RNA polymerase III-dependent transcription through phosphorylation and inhibition of MAF1 an RNA polymerase III-repressor. In parallel to protein synthesis, also regulates lipid synthesis through SREBF1/SREBP1 and LPIN1. To maintain energy homeostasis mTORC1 may also regulate mitochondrial biogenesis through regulation of PPARGC1A. mTORC1 also negatively regulates autophagy through phosphorylation of ULK1. Under nutrient sufficiency, phosphorylates ULK1 at 'Ser-758', disrupting the interaction with AMPK and preventing activation of ULK1. Also prevents autophagy through phosphorylation of the autophagy inhibitor DAP. Also prevents autophagy by phosphorylating RUBCNL/Pacer under nutrient-rich conditions. mTORC1 exerts a feedback control on upstream growth factor signaling that includes phosphorylation and activation of GRB10 a INSR-dependent signaling suppressor. Among other potential targets mTORC1 may phosphorylate CLIP1 and regulate microtubules. As part of the mTORC2 complex MTOR may regulate other</p>

cellular processes including survival and organization of the cytoskeleton. Plays a critical role in the phosphorylation at 'Ser-473' of AKT1, a pro-survival effector of phosphoinositide 3-kinase, facilitating its activation by PDK1. mTORC2 may regulate the actin cytoskeleton, through phosphorylation of PRKCA, PXN and activation of the Rho-type guanine nucleotide exchange factors RHOA and RAC1A or RAC1B. mTORC2 also regulates the phosphorylation of SGK1 at 'Ser-422'. Regulates osteoclastogenesis by adjusting the expression of CEBPB isoforms. Plays an important regulatory role in the circadian clock function. regulates period length and rhythm amplitude of the suprachiasmatic nucleus (SCN) and liver clocks. Phosphorylates SQSTM1, promoting interaction between SQSTM1 and KEAP1 and subsequent inactivation of the BCR(KEAP1) complex.

Cellular Localization

Endoplasmic Reticulum Membrane Peripheral Membrane Protein Cytoplasmic Side Golgi Apparatus Membrane Mitochondrion Outer Membrane Lysosome Cytoplasm Nucleus Pml Body Microsome Membrane Lysosome Membrane Shuttles Between Cytoplasm And Nucleus Accumulates In The Nucleus In Response To Hypoxia (By Similarity) Targeting To Lysosomes Depends On Amino Acid Availability And RragA And RragB Lysosome Targeting Also Depends On Interaction With Meak7 Translocates To The Lysosome Membrane In The Presence Of Tm4sf5

Post-translational Modifications

Autophosphorylates when part of mTORC1 or mTORC2. Phosphorylation at Ser-1261, Ser-2159 and Thr-2164 promotes autophosphorylation. Phosphorylation in the kinase domain modulates the interactions of MTOR with RPTOR and PRAS40 and leads to increased intrinsic mTORC1 kinase activity. Phosphorylation at Thr-2173 in the ATP-binding region by AKT1 strongly reduces kinase activity.